

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

SCIENCE.

EDITORIAL COMMITTEE: S. NEWCOMB, Mathematics; R. S. WOODWARD, Mechanics; E. C. PICKERING, Astronomy; T. C. MENDENHALL, Physics; R. H. Thurston, Engineering; Ira Remsen, Chemistry; J. Le Conte, Geology; W. M. Davis, Physiography; O. C. Marsh, Paleontology; W. K. Brooks, Invertebrate Zoölogy; C. Hart Merriam, Vertebrate Zoölogy; S. H. Scudder, Entomology; N. L. Britton, Botany; Henry F. Osborn, General Biology; H. P. Bowditch, Physiology; J. S. Billings, Hygiene; J. McKeen Cattell, Psychology; Daniel G. Brinton, J. W. Powell, Anthropology.

FRIDAY, JUNE 21, 1895.

CONTENTS:

On the Distribution and the Secular Variation of Terrestrial Magnetism: L. A. BAUER673
On a Devonian Limestone-Breccia in Southwestern Missouri: OSCAR HERSHEY676
Current Notes on Physiography (X.): W. M. DAVIS
Notes on Agriculture (III.): BYRON D. HALSTED 680
Correspondence: —
Scientific Literature:—
Notes and News: —
Societies and Academies:—
Scientific Journals: —

MSS. intended for publication and books, etc., intended for review should be sent to the responsible editor, Prof. J. McKeen Cattell, Garrison on Hudson, N. Y. Subscriptions and advertisements should be sent to SCIENCE, 41 N. Queen St., Lancaster, Pa., or 41 East 49th St., New York.

ON THE DISTRIBUTION AND THE SECULAR VARIATION OF TERRESTRIAL MAGNETISM.

In two papers* read before the Philosophical Society of Washington, May 25th, the following main results were obtained:

* 'On the Secular Variation of Terrestrial Magnetism' and 'A Preliminary Analysis of the Problem of Terrestrial Magnetism and its Variations.'

The minimum change in declination along a parallel of latitude at any particular time, and the minimum average secular change along a parallel of latitude during a given interval of time occur near the equator; both quantities generally increase on leaving the equator.

Exactly the reverse is the case with regard to the inclination, viz.:—

The maximum change in inclination along a parallel of latitude at any particular time, and the maximum average secular change along a parallel of latitude during a given interval of time occur near the equator; both quantities generally diminish on leaving the equator.

These laws were established with the aid of data scaled from magnetic charts from 1780 to 1885 at points 20° distant in longitude and in latitudes 60°N, 40°N, 20°N, equator, 20°S, 40°S and 60°S. They again point to the same conclusion reached previously by the writer in a somewhat different way, namely, that the distribution and the secular variation of terrestrial magnetism appear to be closely related; they are subject to similar laws. It is hence probable that they are both to be referred primarily to the same cause. This common cause seems to be connected in some way with the earth's rotation.

If we regard the earth as uniformly magnetized, having its magnetic poles coincident with the geographical poles, and take the X axis of a system of coördinates whose origin is in the center of the earth, parallel to the magnetic axis, we shall get the fol-